

SEMI-FREDHOLM OPERATORS AND PERTURBATION FUNCTIONS

Snežana Živković–Zlatanović

Abstract. In this paper we give several remarks on [3] and a different proof of the inequalities in [8]. Also we introduce the concept of a lower perturbation function, and prove that some usual measures of noncompactness of operators and also some measures of non-strict-cosingularity of operators are lower perturbation functions. For each lower perturbation function δ , we define functions ∇_δ and K_δ , and give the connection with lower semi-Fredholm operators.

1. Introduction and preliminaries

In this paper X , Y and Z are complex Banach spaces, $B(X, Y)$ (respectively $K(X, Y)$) the set of all bounded (respectively compact) linear operators from X into Y . We shall write $B(X)$ ($K(X)$) instead of $B(X, X)$ ($K(X, X)$).

An operator $T \in B(X, Y)$ is in $\Phi_+(X, Y)$ ($\Phi_-(X, Y)$) if the range $R(T)$ is closed and the dimension $\alpha(T)$ of the null space $N(T)$ of T is finite (the codimension $\beta(T)$ of $R(T)$ in Y is finite). Operators in $\Phi_+(X, Y) \cup \Phi_-(X, Y)$ are called semi-Fredholm operators. For such operators the index, $i(T)$, is defined by $i(T) = \alpha(T) - \beta(T)$. We set $\Phi(X, Y) = \Phi_+(X, Y) \cap \Phi_-(X, Y)$. The operators in $\Phi(X, Y)$ are called Fredholm. We shall write $\Phi_+(X)$ (resp. $\Phi_-(X)$, $\Phi(X)$) instead of $\Phi_+(X, X)$ (resp. $\Phi_-(X, X)$, $\Phi(X, X)$).

Recall that the essential spectral radius of $T \in B(X)$, $r_e(T)$, is defined by

$$r_e(T) = \max\{|\lambda| : T - \lambda I \notin \Phi(X)\}.$$

Let B_X denote the closed unit ball of X . Let $T \in B(X, Y)$ and

$$m(T) = \inf\{\|Tx\| : \|x\| = 1\}$$

be the *minimum modulus* of T , and let

$$n(T) = \sup\{\epsilon \geq 0 : \epsilon B_Y \subset TB_X\}$$

Received May 7, 1997

1991 *Mathematics Subject Classification*: 47A10, 47A53, 47A55.

Supported by Grant 04M03 of RFNS through Math. Inst. SANU

be the *surjection modulus* of T . Recall that $m(T^*) = n(T)$, where $T^* \in B(Y^*, X^*)$ is the adjoint operator.

If M is a subspace of X , then J_M will denote the embedding map of M into X , and if V is a subspace of Y , then Q_V will denote the canonical map of Y onto the quotient space Y/V .

An operator $T \in B(X, Y)$ is *strictly singular* ($T \in S(X, Y)$) if, for every infinite dimensional (closed) subspace M of X , the restriction of T to M , $T|_M$, is not a homeomorphism, i.e. $m(T|_M) = 0$. An operator $T \in B(X, Y)$ is *strictly cosingular* ($T \in CS(X, Y)$) if, for every infinite codimensional closed subspace V of Y the composition $Q_V T$ is not surjective.

For $A \in B(X, Y)$, set

$$\|A\|_C = \inf\{\|A + K\| : K \in K(X, Y)\}.$$

If Ω is a non-empty bounded subset of X , then the Kuratowski measure of noncompactness of Ω is denoted by $\alpha(\Omega)$, and

$$\alpha(\Omega) = \inf\{\epsilon > 0 : \Omega \subset \cup_{i=1}^n D_i, D_i \subset X, \text{diam} D_i \leq \epsilon\}.$$

For $A \in B(X, Y)$ the Kuratowski measure of noncompactness of A is denoted by $\|A\|_\alpha$ and defined by

$$\|A\|_\alpha = \inf\{k \geq 0 : \alpha(A\Omega) \leq k\alpha(\Omega), \Omega \subset X \text{ is bounded}\}.$$

It is easy to see that $\|A\|_\alpha = \sup\{\alpha(A\Omega) : \Omega \subset X, \alpha(\Omega) = 1\}$.

If Ω is a non-empty bounded subset of X , then the Hausdorff measure of noncompactness of Ω , is denoted by $q(\Omega)$, and

$$q(\Omega) = \inf\{\epsilon > 0 : \Omega \text{ has a finite } \epsilon\text{-net in } X\}.$$

For $A \in B(X, Y)$ the Hausdorff measure of noncompactness of A is denoted by $\|A\|_q$ and defined by

$$\|A\|_q = \inf\{k \geq 0 : q_Y(A\Omega) \leq kq_X(\Omega), \Omega \subset X \text{ is bounded}\}.$$

Recall that $\|A\|_q = q_Y(AB_X)$.

For $A \in B(X, Y)$, set (see [6])

$$\|A\|_\mu = \inf\{\|A|_L\| : L \text{ subspace of } X, \text{codim} L < \infty\},$$

and

$$\begin{aligned} \Gamma_M(A) &= \inf_{N \subset M} \|A|_N\|, \quad \Gamma(A) = \Gamma_X(A), \\ \Delta_M(A) &= \sup_{N \subset M} \Gamma_N(A), \quad \Delta(A) = \Delta_X(A), \end{aligned}$$

where M, N denote infinite dimensional subspaces of X (see [9]). Schechter [9] proved that Δ is a submultiplicative seminorm and

$$(1.1) \quad \Delta(A) = 0 \iff A \in S(X, Y).$$

For $A \in B(X, Y)$, set (see [12])

$$\begin{aligned} K_V(A) &= \inf_{W \supset V} \|Q_W A\|, & K(A) &= K_{\{0\}}(A), \\ \nabla_V(A) &= \sup_{W \supset V} K_W(A), & \nabla(A) &= \nabla_{\{0\}}(A), \end{aligned}$$

where V, W denote closed infinite codimensional subspaces of Y .

Recall that [12],

$$(1.2) \quad \begin{aligned} K(A) > 0 &\iff A \in \Phi_-(X, Y), \\ \nabla(A) = 0 &\iff A \in CS(X, Y). \end{aligned}$$

Zemánek [13] considered the following functions

$$\begin{aligned} u(A) &= \sup\{m(A|_W) : W \text{ is closed subspace of } X \text{ with } \dim W = \infty\}, \\ v(A) &= \sup\{n(Q_V A) : V \text{ is closed subspace of } Y \text{ with } \operatorname{codim} V = \infty\}. \end{aligned}$$

From the definition of the strictly singular and strictly cosingular operators it is obvious that

$$(1.3) \quad u(A) = 0 \iff A \in S(X, Y),$$

$$(1.4) \quad v(A) = 0 \iff A \in CS(X, Y).$$

2. Results

Recall that F. Galaz-Fontes [3] introduced and investigated a perturbation function.

Definition 2.1. A perturbation function is a function γ , assigning to each pair of complex Banach spaces X, Y , and $T \in B(X, Y)$ a nonnegative number $\gamma(T)$, with the following properties:

$$(2.1.1) \quad \gamma(\lambda T) = |\lambda| \gamma(T), \quad \lambda \in \mathbb{C},$$

$$(2.1.2) \quad \gamma(T + K) = \gamma(T), \quad K \in K(X, Y),$$

$$(2.1.3) \quad \gamma(T) \leq \|T\|,$$

$$(2.1.4) \quad m(T) \leq \gamma(T),$$

$$(2.1.5) \quad \gamma(T|_M) \leq \gamma(T),$$

where M denotes an infinite dimensional subspace of X .

He proved that the quantities $\|\cdot\|_C$, $\|\cdot\|_\alpha$, $\|\cdot\|_q$, $\|\cdot\|_\mu$, Δ and u are perturbation functions.

Let us note that the following quantity is a measure of non-strict-singularity and we shall show that it is a perturbation function.

Example 2.2. For $T \in B(X, Y)$, set

$$\|T\|_S = \inf\{\|T + S\| : S \in S(X, Y)\}.$$

(2.1.1)-(2.1.3) follow easily from the definition.

Since Δ is a seminorm which annihilates precisely on the strictly singular operators [9] and such that $\Delta(T) \leq \|T\|$, $T \in B(X, Y)$, it follows that

$$\Delta(T) = \Delta(T + S) \leq \|T + S\|, \text{ for each } S \in S(X, Y).$$

Therefore

$$\Delta(T) \leq \|T\|_S.$$

Since $m(T) \leq u(T) \leq \Delta(T)$, we get $m(T) \leq \|T\|_S$. (2.1.5) follows from the fact that a restriction of a strictly singular operator to an infinite dimensional subspace of X is a strictly singular operators also.

Recall that for a given perturbation function γ , and $T \in B(X, Y)$, F. Galaz-Fontes [3] defined

$$\begin{aligned} \Gamma_{\gamma, M}(T) &= \inf\{\gamma(T|_V) : V \subset M\}, \quad \Gamma_\gamma(T) = \Gamma_{\gamma, X}(T); \\ \Delta_{\gamma, M}(T) &= \sup\{\Gamma_{\gamma, V}(T) : V \subset M\}, \quad \Delta_\gamma(T) = \Delta_{\gamma, X}(T), \end{aligned}$$

where M, V denote closed infinite dimensional subspaces of X .

In the following γ is a perturbation function. F. Galaz-Fontes proved that if $T \in S(X, Y)$, then $\Delta_\gamma(T) = 0$ [3, Proposition 9]. Actually, we shall prove that the equivalence holds.

Proposition 2.3. *Let $T \in B(X, Y)$. Then*

$$(2.3.1) \quad T \in S(X, Y) \iff \Delta_\gamma(T) = 0.$$

Proof. First we show:

$$(2.3.2) \quad u(T) \leq \Delta_\gamma(T) \leq \Delta(T).$$

From (2.1.3) and the definitions of Δ and Δ_γ it follows that $\Delta_\gamma(T) \leq \Delta(T)$.

Let M be a closed infinite dimensional subspace of X . We have

$$\gamma(T|_V) \geq m(T|_V) \geq m(T|_M),$$

for each closed infinite dimensional subspace V of M . It implies

$$\Gamma_{\gamma,M}(T) = \inf_{V \subset M} \gamma(T|_V) \geq m(T|_M).$$

Hence

$$\Delta_\gamma(T) = \sup_M \Gamma_{\gamma,M}(T) \geq \sup_M m(T|_M) = u(T).$$

(2.3.1) follows directly from (2.3.2), (1.1) and (1.3). $\square$

Let us remark that if γ is a submultiplicative seminorm then it can be proved (analogously as in [9]) that Δ_γ is a submultiplicative seminorm and

$$\Gamma_\gamma(T + S) \leq \Gamma_\gamma(T) + \Delta_\gamma(S), \quad T, S \in B(X, Y).$$

If $\gamma = \|\cdot\|_q$ we shall write Γ_q (Δ_q) instead of Γ_γ (Δ_γ). Analogously, we introduce Γ_α (Δ_α), Γ_μ (Δ_μ), Γ_C (Δ_C), Γ_u (Δ_u) and Γ_Δ (Δ_Δ). By [3, Lemma 8] it follows that $\Gamma_\Delta = \Gamma$ and $\Delta_\Delta = \Delta$.

Lemma 2.4. *If there exists a constant $c > 0$ such that $\gamma(T) \geq c\Delta(T)$, for each $T \in B(X, Y)$, then*

$$(2.4.1) \quad \Gamma_\gamma(T) \leq \Gamma(T) \leq c^{-1}\Gamma_\gamma(T),$$

$$(2.4.2) \quad \Delta_\gamma(T) \leq \Delta(T) \leq c^{-1}\Delta_\gamma(T), \quad T \in B(X, Y).$$

Proof. (2.4.1). From (2.1.3) and the definitions of Γ and Γ_γ it follows that $\Gamma_\gamma(T) \leq \Gamma(T)$. Further from the hypothesis and [3, Lemma 8] we get

$$\Gamma_\gamma(T) \geq c\Gamma_\Delta(T) = c\Gamma(T).$$

(2.4.2) can be proved analogously to (2.4.1). $\square$

Recall that

$$\begin{aligned} \|T\|_\mu &\geq \Delta(T) \text{ [9, Theorem 2.10]}, \\ \|T\|_q &\geq 2^{-1}\|T\|_\mu \geq 2^{-1}\Delta(T) \text{ [6, Theorem 3.1]}, \\ \|T\|_\alpha &\geq 2^{-1}\|T\|_q \geq 4^{-1}\Delta(T), \\ \|T\|_C &\geq \|T\|_\mu \geq \Delta(T), \\ \|T\|_S &\geq \Delta(T), \quad T \in B(X, Y). \end{aligned}$$

By Lemma 2.4 we get

$$\begin{aligned} \Gamma_\mu(T) &= \Gamma(T), \quad \Delta_\mu(T) = \Delta(T), \\ \Gamma_q(T) &\leq \Gamma(T) \leq 2\Gamma_q(T), \quad \Delta_q(T) \leq \Delta(T) \leq 2\Delta_q(T), \\ (2.5) \quad \Gamma_\alpha(T) &\leq \Gamma(T) \leq 4\Gamma_\alpha(T), \quad \Delta_\alpha(T) \leq \Delta(T) \leq 4\Delta_\alpha(T), \\ \Gamma_C(T) &= \Gamma(T), \quad \Delta_C(T) = \Delta(T), \\ \Gamma_S(T) &= \Gamma(T), \quad \Delta_S(T) = \Delta(T). \end{aligned}$$

The inequalities in (2.5) were proved in [8] and [5, Proposition 3.9]. However, our proof is different from the proofs in [8] and [5].

Remark 2.6. Clearly, $\Delta_\gamma \leq \gamma$, and by (2.3.2), for $\gamma = u$, we get

$$\Delta_u = u.$$

Recall that [3, Lemma 11]

$$B \leq \Gamma_u \leq \Gamma,$$

where $B(T) = \sup\{m(T|_V) : V \text{ closed subspace of } X, \text{codim} V < \infty\}$. M. González and A. Martínón proved that the quantities B and Γ_u are not equivalent and also that the quantities Γ_u and Γ are not equivalent [5, Theorem 2.7 and Corollary 3.5].

In the terminology of [10], by [3, Theorem 6] it results that u/Γ_u is a perturbation function for $\Phi_+(X, Y)$. Since $\Gamma_u \geq B$ it follows that the perturbation function u/Γ_u is better than the perturbation function u/B . Thus, Theorem 2.14 in [9] is a consequence of the fact that u/Γ_u is a perturbation function for $\Phi_+(X, Y)$.

The above fact can be established from the inequality [7, Proposición 25.8.4]

$$\Gamma_u(T + S) \leq u(T) + \Gamma(S), \quad T, S \in B(X, Y).$$

Indeed, let $u(S) < \Gamma_u(T)$. Then

$$\Gamma_u(T) = \Gamma_u(T + S + (-S)) \leq \Gamma(T + S) + u(S) < \Gamma(T + S) + \Gamma_u(T),$$

and

$$\Gamma(T + S) > 0 \implies T + S \in \Phi_+(X, Y).$$

Now we introduce and investigate a new function connected with the lower semi-Fredholm operators.

Definition 2.7. A lower perturbation function is a function δ , which assigns to each pair of complex Banach spaces X, Y , and $T \in B(X, Y)$ a nonnegative number $\delta(T)$, with the following properties:

$$(2.7.1) \quad \delta(\lambda T) = |\lambda| \delta(T),$$

$$(2.7.2) \quad \delta(T + K) = \delta(T), \quad K \in K(X, Y),$$

$$(2.7.3) \quad \delta(T) \leq \|T\|,$$

$$(2.7.4) \quad n(T) \leq \delta(T),$$

$$(2.7.5) \quad U \subset V \implies \delta(Q_U T) \geq \delta(Q_V T),$$

where U, V denote closed infinite codimensional subspaces of Y .

We shall give several examples.

Example 2.8. The quantity $\|\cdot\|_\mu$ is a lower perturbation function.

We shall prove only (2.7.4) and (2.7.5). Recall that $\|T\|_\mu = \|T^*\|_q$ [1, Teorema 2.5.2]. Now from [3, Example 3] it follows:

$$\|T\|_\mu = \|T^*\|_q \geq m(T^*) = n(T).$$

Let U and V denote closed infinite codimensional subspaces of Y and $U \subset V$. Let be $\phi : Y/U \rightarrow Y/V$ a map defined by $\phi(y + U) = y + V$. Then, by [6, Lemma 3.2] we have

$$\|Q_V T\|_\mu = \|\phi Q_U T\|_\mu \leq \|\phi\|_\mu \|Q_U T\|_\mu \leq \|\phi\| \|Q_U T\|_\mu \leq \|Q_U T\|_\mu.$$

Example 2.9. The quantity $\|\cdot\|_q$ is a lower perturbation function.

We shall prove only (2.7.4) and (2.7.5). Recall that

$$\|T\|_q \geq \|T^*\|_\mu$$

(see [4, Theorem 1 (ii), Proposition 6 (ii)] or [1, Corollary 2.5.4]). Since $\|T^*\|_\mu \geq m(T^*) = n(T)$, we get

$$\|T\|_q \geq n(T).$$

Analogously as in Example 2.8, it can be proved that $\|\cdot\|_q$ has the property (2.7.5).

Example 2.10. The quantity $\|\cdot\|_C$ is a lower perturbation function.

Indeed, since $\|T\|_C \geq \|T\|_q$, we get $\|T\|_C \geq n(T)$. Since $B(Y, Z)K(X, Y) \subset K(X, Z)$, it follows that $\|BA\|_C \leq \|B\| \|A\|_C$, $A \in B(X, Y)$, $B \in B(Y, Z)$. In an analogous way as in Example 2.8, we obtain (2.7.5).

Example 2.11. The quantity ∇ is a lower perturbation function.

Since ∇ is a seminorm on $B(X, Y)$ which annihilates precisely on the set $CS(X, Y)$ and $K(X, Y) \subset CS(X, Y)$ we obtain properties (2.7.1) and (2.7.2). (2.7.3) is obvious.

To prove the property (2.7.4), suppose that V is a closed subspace of Y with $\text{codim} V \geq 1$. Then V° is a subspace of Y^* , $\dim V^\circ = \text{codim} V \geq 1$ and $\|Q_V T\| = \|T^* J_{V^\circ}\|$. Thus

$$\begin{aligned} & \{\|Q_V T\| : V \text{ closed subspace of } Y, \text{codim} V \geq 1\} \\ & \subset \{\|T^* J_M\| : M \text{ subspace of } Y^*, \dim M \geq 1\}. \end{aligned}$$

Hence

$$\begin{aligned} \nabla(T) & \geq K(T) = \inf\{\|Q_V T\| : V \text{ closed subspace of } Y, \text{codim} V = \infty\} \\ & \geq \inf\{\|Q_V T\| : V \text{ closed subspace of } Y, \text{codim} V \geq 1\} \\ & \geq \inf\{\|T^* J_M\| : M \text{ subspace of } Y^*, \dim M \geq 1\} \\ & = \inf\{\|T^* J_M\| : M \text{ subspace of } Y^*, \dim M = 1\} \\ & = \inf\{\|T^* f\| : f \in Y^*, \|f\| = 1\} = m(T^*) = n(T). \end{aligned}$$

Let us remark that

$$n(T) \leq K(T) \leq \nabla(T) \leq \|T\|_q.$$

Since $\nabla(BA) \leq \nabla(B)\nabla(A)$, $A \in B(X, Y)$, $B \in B(Y, Z)$, in an analogous way as in Example 2.8, we obtain the property (2.7.5).

Example 2.12. For $T \in B(X, Y)$, set

$$\|T\|_{CS} = \inf\{\|T + C\| : C \in CS(X, Y)\}.$$

The quantity $\|\cdot\|_{CS}$ is a lower perturbation function.

Really, as in the previous example, we obtain that $\|\cdot\|_{CS}$ has properties (2.7.1)-(2.7.3). It is easy to see that $\nabla(T) \leq \|T\|_{CS}$ and since $n(T) \leq \nabla(T)$, we get $n(T) \leq \|T\|_{CS}$. Since $B(Y, Z)CS(X, Y) \subset CS(X, Z)$, it follows that $\|BA\|_{CS} \leq \|B\|\|A\|_{CS}$, $A \in B(X, Y)$, $B \in B(Y, Z)$. Now, in an analogous way as in Example 2.8, we obtain the property (2.7.5).

Problem: The quantity v has properties (2.7.1)-(2.7.4), but we do not know whether it has property (2.7.5).

From Definition 2.7 we have

Lemma 2.13. *If δ is a lower perturbation function, then:*

$$(2.13.1) \quad \delta(K) = 0, \quad K \in K(X, Y),$$

$$(2.13.2) \quad \delta(Q_V) = 1,$$

V closed infinite codimensional subspace of Y .

Proof. (2.13.2) follows from the following inequalities:

$$1 = m(J_{V^\circ}) = n(Q_V) \leq \delta(Q_V) \leq \|Q_V\| = 1. \quad \square$$

In the following δ is a lower perturbation function.

Lemma 2.14. *Let $P \in B(X)$. If $\delta(P) < 1$, then $I + P \in \Phi(X)$ and $i(I + P) = 0$.*

Proof. Assume that $I + P \notin \Phi_-(X)$. By [6, Lemma 5.4] there exists $K \in K(X)$ such that $\text{codim } \overline{R(I + P - K)} = \infty$. Set $U = \overline{R(I + P - K)}$. From $Q_U(I + P - K) = 0$ we get $Q_U = Q_U(K - P)$. Hence, by (2.13.2), (2.7.2) and (2.7.5), it follows that

$$1 = \delta(Q_U) = \delta(Q_U(K - P)) = \delta(Q_U P) \leq \delta(P).$$

This contradicts the hypothesis. Thus, $I + P \in \Phi_-(X)$.

Let $0 \leq \lambda \leq 1$. Then $\delta(\lambda P) < 1$ and therefore $I + \lambda P \in \Phi_-(X)$. Since the index is locally constant it follows that $i(I + P) = i(I) = 0$. Consequently, $I + P \in \Phi(X)$. $\square$

The following proposition can be proved (see [3, Theorem 5]).

Proposition 2.15. $r_e(T) = \lim_{n \rightarrow \infty} (\delta(T^n))^{\frac{1}{n}}, \quad T \in B(X).$

Definition 2.16. For $T \in B(X, Y)$, set

$$\begin{aligned} K_{\delta, V}(T) &= \inf \{ \delta(Q_W T) : W \supset V \}, \quad K_\delta(T) = K_{\delta, \{0\}}(T), \\ \nabla_{\delta, V}(T) &= \sup \{ K_{\delta, W}(T) : W \supset V \}, \quad \nabla_\delta(T) = \nabla_{\delta, \{0\}}(T), \end{aligned}$$

where V, W denote closed infinite codimensional subspaces of Y .

Theorem 2.17. Let $S, T \in B(X, Y)$. If $\nabla_\delta(T) < K_\delta(T)$, then $T, T + S \in \Phi_-(X, Y)$ and $i(T + S) = i(T)$.

Proof. Suppose that $T + S \notin \Phi_-(X, Y)$. Then, by [6, Lemma 5.4], there is $K \in K(X, Y)$ such that $\text{codim} R(T + S - K) = \infty$. Set $U = \overline{R(T + S - K)}$. Let V be a closed infinite codimensional subspace of Y such that $V \supset U$. Then $Q_V(T + S - K) = 0$, and $Q_V T = Q_V(K - S)$. Therefore

$$\begin{aligned} K_\delta(T) &\leq K_{\delta, U}(T) = \inf \{ \delta(Q_V T) : V \supset U \} = \inf \{ \delta(Q_V(K - S)) : V \supset U \} \\ &= \inf \{ \delta(Q_V S) : V \supset U \} = K_{\delta, U}(S) \leq \nabla_\delta(S). \end{aligned}$$

This contradicts the hypothesis.

Let $0 \leq \lambda \leq 1$. Then $\nabla_\delta(\lambda S) < K_\delta(T)$. It follows that $T + \lambda S \in \Phi_-(X, Y)$. Thus $T, T + S \in \Phi_-(X, Y)$ and, since the index is locally constant, we get $i(T) = i(T + S)$. $\square$

Theorem 2.18. Let $T \in B(X, Y)$. Then $T \in \Phi_-(X, Y) \iff K_\delta(T) > 0$.

Proof. Let $K_\delta(T) > 0$. If $S = 0$ then $\nabla_\delta(S) = 0 < K_\delta(T)$, and by Theorem 2.17 we get $T \in \Phi_-(X, Y)$.

Let $T \in \Phi_-(X, Y)$. Then $\text{codim} R(T) < \infty$, and we can express Y as a direct sum $Y = R(T) \oplus V$ where V is a subspace of Y with $\dim V < \infty$. This implies that $Q_V T$ is surjective, i.e. $n(Q_V T) > 0$. Let W be a closed infinite codimensional subspace of Y . Clearly, $\text{codim}(V + W) = \infty$. Now, by (2.7.4) and (2.7.5), we have:

$$n(Q_V T) \leq n(Q_{V+W} T) \leq \delta(Q_{V+W} T) \leq \delta(Q_W T).$$

Consequently,

$$\begin{aligned} K_\delta(T) &= \inf \{ \delta(Q_W T) : W \text{ closed subspace of } Y, \text{ codim } W = \infty \} \\ &\geq n(Q_V T) > 0. \quad \square \end{aligned}$$

Proposition 2.19. $T \in B(X, Y)$ is strictly cosingular if and only if $\nabla_\delta(T) = 0$.

Proof. First we shall prove the following inequality

$$(2.19.1) \quad v(T) \leq \nabla_\delta(T) \leq \nabla(T).$$

The right side of the above inequality follows from (2.7.3) and the definitions of ∇ and ∇_δ . To prove the left side, let V be a closed infinite codimensional subspace of Y . We have

$$\delta(Q_W T) \geq n(Q_W T) \geq n(Q_V T),$$

for each closed infinite codimensional subspace W , such that $W \supset V$. It implies

$$K_{\delta, V}(T) = \inf_{W \supset V} \delta(Q_W T) \geq n(Q_V T).$$

Hence

$$\nabla_\delta(T) = \sup_V K_{\delta, V}(T) \geq \sup_V n(Q_V T) = v(T).$$

Now the assertion of Proposition follows from (2.19.1), (1.2) and (1.4). $\square$

Let us remark that if δ is a submultiplicative seminorm then it can be proved that ∇_δ is a submultiplicative seminorm and

$$K_\delta(T + S) \leq K_\delta(T) + \nabla_\delta(S), \quad T, S \in B(X, Y).$$

Also in this case we can show that ∇_δ is a lower perturbation function with $\nabla_\delta(T) \leq \delta(T)$ (the property (2.7.4) follows from the inequality $n(T) \leq v(T) \leq \nabla_\delta(T)$ and the property (2.7.5) can be proved analogously as in Example 2.8). Since

$$M(T) \leq K_\delta(T) \leq K(T),$$

where $M(T) = \sup\{n(Q_V T) : \dim V < \infty\}$ [13], from [13, Theorem 8.1] it follows:

$$s_-(T) = \lim_{n \rightarrow \infty} (K_\delta(T^n))^{\frac{1}{n}},$$

where $s_-(T) = \inf\{|\lambda| : \lambda I - T \notin \Phi_-(X)\}$.

The next lemma implies that $\nabla(\nabla_\delta) = \nabla_\delta$.

Lemma 2.20. $K_{\delta, V}(T) = \inf\{\nabla_{\delta, W}(T) : W \supset V\}$, where V, W denote closed infinite codimensional subspaces of Y .

Proof. Since

$$\nabla_{\delta, W}(T) \geq K_{\delta, W}(T) \geq K_{\delta, V}(T),$$

for each W with $W \supset V$, we get

$$\inf\{\nabla_{\delta, W}(T) : W \supset V\} \geq K_{\delta, V}(T).$$

In the following M, N are closed infinite codimensional subspace of Y . Let $M \supset W$. From (2.7.5) it follows that

$$K_{\delta, M}(T) = \inf_{N \supset M} \delta(Q_N T) \leq \delta(Q_M T) \leq \delta(Q_W T).$$

Consequently

$$\nabla_{\delta, W}(T) = \sup_{M \supset W} K_{\delta, M}(T) \leq \delta(Q_W T).$$

This implies

$$\inf_{W \supset V} \nabla_{\delta, W}(T) \leq \inf_{W \supset V} \delta(Q_W T) = K_{\delta, V}(T). \quad \square$$

If $\delta = \|\cdot\|_q$ we shall write $K_q(\nabla_q)$ instead of $K_\delta(\nabla_\delta)$. Analogously, we introduce $K_\mu(\nabla_\mu)$, $K_C(\nabla_C)$, $K_\nabla(\nabla_\nabla)$ i $K_{CS}(\nabla_{CS})$. From Lemma 2.20 it follows that $K_\nabla = K$ and $\nabla_\nabla = \nabla$.

In an analogous way as Lemma 2.4 the next lemma can be proved.

Lemma 2.21. *If there exists a constant $c > 0$ such that $\delta(T) \geq c\nabla(T)$, for each $T \in B(X, Y)$, then*

$$\begin{aligned} K_\delta(T) &\leq K(T) \leq \frac{1}{c} K_\delta(T), \\ \nabla_\delta(T) &\leq \nabla(T) \leq \frac{1}{c} \nabla_\delta(T), \quad T \in B(X, Y). \end{aligned}$$

Recall that

$$\begin{aligned} \|T\|_q &\geq \nabla(T), \\ \|T\|_\mu &\geq \frac{1}{2} \|T\|_q \geq \frac{1}{2} \nabla(T), \\ \|T\|_C &\geq \|T\|_q \geq \nabla(T), \\ \|T\|_{CS} &\geq \nabla(T), \quad T \in B(X, Y). \end{aligned}$$

Now, by Lemma 2.21 we obtain

$$\begin{aligned} (2.22) \quad K_q(T) &= K(T), \quad \nabla_q(T) = \nabla(T), \\ K_\mu(T) &\leq K(T) \leq 2K_\mu(T), \quad \nabla_\mu(T) \leq \nabla(T) \leq 2\nabla_\mu(T), \\ K_C(T) &= K(T), \quad \nabla_C(T) = \nabla(T), \\ K_{CS}(T) &= K(T), \quad \nabla_{CS}(T) = \nabla(T). \end{aligned}$$

The equalities in (2.22) were proved in [11, Summary and discussion, Remark 2]. However, our proof is different from this one.

Acknowledgment. I am grateful to Professor Vladimir Rakočević for helpful comments and suggestions.

References

- [1] P. P. Ахмеров, М. И. Каменский, А. С. Потапов и др., *Меры некомпактности и уплотняющие операторы*, Наука, Новосибирск, 1986.
- [2] A. S. Fainstein, *On measures of noncompactness of linear operators and analogous of the minimum modulus for semi-Fredholm operators (Russian)*, Spektral'naja Teorija Operatorov i Prilozh., Baku **6** (1985), 182–195..
- [3] F. Galaz-Fontes, *Measures of noncompactness and upper semi-Fredholm perturbation theorems*, Proc. Amer. Math. Soc. **118** (1993), 891–897.
- [4] M. González and A. Martínón, *Operational quantities derived from the norm and measures of non-compactness*, Proc. R. Ir. Acad. **91A** (1991), 63–70.
- [5] M. González and A. Martínón, *Operational quantities characterizing semi-Fredholm operators*, Studia Math. **114** (1995), 13–27.
- [6] A. Lebow and M. Schechter, *Semigroups of operators and measures of noncompactness*, J. Funct. Anal. **7** (1971), 1–26.
- [7] A. Martínón, *Cantidades operacionales en teoria de Fredholm*, Doctoral thesis, University of La Laguna, 1989.
- [8] V. Rakočević, *Measures of non-strict singularity of operators*, Mat. Vesnik **35** (1983), 79–82.
- [9] M. Schechter, *Quantities related to strictly singular operators*, Indiana Univ. Math. J. **21** (11) (1972), 1061–1071.
- [10] M. Schechter and R. Whitley, *Best Fredholm perturbation theorems*, Studia Math. **90** (1988).
- [11] H.-O. Tylli, *On semifredholm operators, Calkin algebras and some related quantities*, Dissertation, Helsinki, 1986.
- [12] L. Weist, *Über strikt singuläre und strikt cosinguläre Operatoren in Banachräumen*, Dissertation, Bonn, 1974.
- [13] J. Zemánek, *Geometric characteristics of semi-Fredholm operators and asymptotic behaviour*, Studia Math. **80** (1984), 219–234.

UNIVERSITY OF NIŠ, FACULTY OF PHILOSOPHY, DEPARTMENT OF MATHEMATICS, ĆIRILA
AND METODIJA 2, 18000 NIŠ, YUGOSLAVIJA, SERBIA